

Data release of short baseline ν_e disappearance analysis at ND280 with the run 1-4 data set (2014), arXiv:1410.8811.

First the best-fit χ^2 is given. It corresponds to the χ^2 value in the global minimum. The file contains the $\Delta\chi^2$ for each point of the grid 51×41 , Δm_{eff}^2 (row) Vs $\sin^2 2\theta_{ee}$ (column) with the respect to the global minimum.

$\Delta m_{eff}^2 (eV^2)$:

0., 0.5, 1, 1.5, 2, 2.5, 3, 3.5, 4, 5, 6, 7, 8, 9, 10, 12.5, 15, 17.5, 20, 22.5, 25, 27.5, 30, 32.5, 35, 37.5, 40, 42.5, 45, 47.5, 50, 52.5, 55, 57.5, 60, 62.5, 65, 67.5, 70, 72.5, 75, 77.5, 80, 82.5, 85, 87.5, 90, 92.5, 95, 97.5, 100

$\sin^2 2\theta_{ee}$:

0., 0.025, 0.05, 0.075, 0.1, 0.125, 0.15, 0.175, 0.2, 0.225, 0.25, 0.275, 0.3, 0.325, 0.35, 0.375, 0.4, 0.425, 0.45, 0.475, 0.5, 0.525, 0.55, 0.575, 0.6, 0.625, 0.65, 0.675, 0.7, 0.725, 0.75, 0.775, 0.8, 0.825, 0.85, 0.875, 0.9, 0.925, 0.95, 0.975, 1